

# Jasmonates act with salicylic acid to confer basal thermotolerance in *Arabidopsis thaliana*

Shannon M. Clarke<sup>1,5</sup>, Simona M. Cristescu<sup>2</sup>, Otto Miersch<sup>3</sup>, Frans J. M. Harren<sup>2</sup>, Claus Wasternack<sup>3</sup> and Luis A. J. Mur<sup>4</sup>

<sup>1</sup>Biochemistry Department, University of Otago, PO Box 56, Dunedin, New Zealand; <sup>2</sup>Life Science Trace Gas Exchange Facility, Department of Molecular and Laser Physics, Radboud University, Heyendaalseweg 135, NL 6525 AJ Nijmegen, The Netherlands; <sup>3</sup>Leibniz-Institute of Plant Biochemistry, Weinberg 3, D-06120 Halle/Saale, Germany; <sup>4</sup>Institute of Biological Sciences, University of Aberystwyth, SY23 3DA, UK; <sup>5</sup>Present address: AgResearch Limited, Invermay Agricultural Centre, Puddle Alley, Private Bag 50034, Mosgiel, New Zealand

## Summary

Author for correspondence:

Shannon M. Clarke

Tel: +64 3 4899114

Fax: +64 3 4893739

Email: shannon.clarke@agresearch.co.nz

New Phytologist (2009)

doi: 10.1111/j.1469-8137.2008.02735.x

**Key words:** *Arabidopsis*, *COI1*, *cpr5-1*, *EIN2*, jasmonic acid (JA), *JAR1*, salicylic acid (SA), thermotolerance.

- The *cpr5-1 Arabidopsis thaliana* mutant exhibits constitutive activation of salicylic acid (SA), jasmonic acid (JA) and ethylene (ET) signalling pathways and displays enhanced tolerance of heat stress (HS).
- *cpr5-1* crossed with *jar1-1* (a JA-amino acid synthetase) was compromised in basal thermotolerance, as were the mutants *opr3* (mutated in OPDA reductase3) and *coi1-1* (affected in an E3 ubiquitin ligase F-box; a key JA-signalling component). In addition, heating wild-type *Arabidopsis* led to the accumulation of a range of jasmonates: JA, 12-oxophytodienoic acid (OPDA) and a JA-isoleucine (JA-Ile) conjugate. Exogenous application of methyl jasmonate protected wild-type *Arabidopsis* from HS.
- Ethylene was rapidly produced during HS, with levels being modulated by both JA and SA. By contrast, the ethylene mutant *ein2-1* conferred greater thermotolerance.
- These data suggest that JA acts with SA, conferring basal thermotolerance while ET may act to promote cell death.

## Introduction

Plants are sessile organisms and their ability to cope with exposure to temperatures above optimal growth conditions is crucial for survival. Plants have a natural capacity to ameliorate the effects of heat shock (HS) via a 'basal' thermotolerance mechanism (Hong & Vierling, 2000). In addition, lesser increases in temperature can acclimatize plants against HS via a process known as 'acquired' thermotolerance. The production of heat shock proteins (HSPs) plays an important role in protecting proteins against irreversible heat-induced damage by preventing denaturation and facilitating the refolding of damaged proteins (Boston *et al.*, 1996; Hong & Vierling, 2000; Hong *et al.*, 2003). In addition, a number of signalling pathways have been implicated in thermotolerance including redox, hormonal and calcium-based (Foyer *et al.*, 1997; Dat *et al.*, 1998a,b; Larkindale & Knight, 2002; Pastori & Foyer, 2002; Clarke *et al.*, 2004; Larkindale *et al.*, 2005).

Our studies using *Arabidopsis thaliana* mutants indicate that salicylic acid (SA)-signalling pathways, which were originally defined as being involved in antipathogenic responses and in

systemic-acquired resistance (SAR; Dong, 2001) also promote basal thermotolerance but were dispensable for the acquired form (Clarke *et al.*, 2004). Heat shock was found to induce SA-regulated pathogenesis-related 1 (*PR1*) transcripts and the ability of the nonexpresser of PR1 protein (*npr1-1*) mutant, which is involved in SA-signal transduction, to recover from heat stress was impaired. Correspondingly, the constitutive expresser of PR1 protein (*cpr5-1*) mutant displayed an enhanced basal thermotolerant phenotype. Although the deduced CPR5 sequence revealed no significant homology to any other gene, it has features of a signal transduction protein with a putative nuclear localization signal and five putative transmembrane domains (Clarke *et al.*, 2000).

In addition to activation of the SA pathway in *cpr5-1*, the jasmonic acid (JA)-inducible genes *PDF1.2* and *THI2.1* were also found to be constitutively expressed (Bowling *et al.*, 1997; Clarke *et al.*, 1998). It may therefore be that the enhanced thermotolerant phenotype observed in the *cpr5-1* mutant also involves the JA-signalling pathway. Jasmonates regulate or coregulate a wide range of developmental processes in plants and responses to biotic and abiotic stresses (Wasternack,

2006, 2007; Balbi & Devoto, 2008). Wounding and pathogen infection or exposure to ozone cause endogenous JA accumulation and the expression of JA-responsive genes, for example, proteinase inhibitors and antifungal proteins such as thionins and defensins (Vijayan, *et al.*, 1998; Rao *et al.*, 2000; Kanna *et al.*, 2003; Howe, 2004; Wasternack, 2006). Defects in JA biosynthesis or signalling pathways have been shown to increase the sensitivity to stresses such as pathogen infection or ozone exposure (McConn *et al.*, 1997; Thomma *et al.*, 1998; Overmyer *et al.*, 2000; Stintzi, *et al.*, 2001; Kanna *et al.*, 2003).

The *coi1-1 Arabidopsis* mutant is defective in most of the known JA-dependent responses, indicating that COI1 has a central role in JA signalling (Xie *et al.*, 1998; Reymond *et al.*, 2000, 2004). COI1 is an E3 ubiquitin ligase which tags proteins with ubiquitin as a signal for their proteolytic destruction in the proteasome. There are also several other *Arabidopsis* mutants that show reduced sensitivity to JA, although they are not defective in all the JA responses; for example, the mutants of *jar1/jin4/jai2* allelic series (encoding a JA-amino acid synthetase; Staswick *et al.*, 1992, 2002, Berger *et al.*, 1996; Lorenzo *et al.*, 2004; Staswick & Tiryaki, 2004) and *jin1/jai1* alleles (encoding a nuclear-localised bHLHzip transcription factor, AtMYC2; Lorenzo *et al.*, 2004). Recent work has provided a more comprehensive view of the JA signalling cascade (Katsir *et al.*, 2008; Staswick, 2008). COI1 has been shown to target a class of putative repressor proteins known as JAZ (jasmonate ZIM-domain), one of which interacts with AtMYC2. The COI1–JAZ complex is stabilized by interaction with JA-isoleucine (JA-Ile), the product of JAR1 (Chini *et al.*, 2007; Thines *et al.*, 2007; Yan *et al.*, 2007; Katsir *et al.*, 2008).

The relationship between the JA- and SA- signalling has often been shown to be antagonistic. This antagonism may reflect SA-mediated suppression of JA biosynthesis (Pěna-Cortes *et al.*, 1993; Doares *et al.*, 1995) by reducing the expression of genes encoding JA-biosynthetic enzymes (Laudert & Weiler, 1998; Spoel *et al.*, 2003). However, *Arabidopsis* mutants *cpr5*, *cpr6*, *cet* (constitutive expression of thionin), *cpr22* and *hrl1* (hypersensitive response-like lesions I) (Clarke *et al.*, 1998; Hilpert *et al.*, 2001; Yoshioka *et al.*, 2001; Devadas *et al.*, 2002) show simultaneous expression of SA- and JA- marker genes. Recently, Mur *et al.* (2006) suggested that the reciprocal SA and JA antagonism of gene expression was dose dependent.

In *cpr5*, JA-mediated effects on plant pathogen resistance were influenced by ethylene (ET, Clarke *et al.*, 2000). Similarly, resistance to necrotrophic pathogens (Penninckx *et al.*, 1998) and CaMV (Love *et al.*, 2005) involved ET–JA interactions. In some necrotrophic interactions, for example, between *Cochliobolus victoriae* and oats (Navarre & Wolpert, 1999), ET will contribute to the elicitation of cell death. However, within the context of thermotolerance, ET may act mainly in basal thermotolerance as this was increased on treatment with the ET precursor 1-aminocyclopropane-1-carboxylic acid (ACC; Larkindale & Knight, 2002). Further, basal rather than acquired thermotolerance was most compromised in ET

receptor mutant *etr1* and in *ein2-1*, which is mutated within membrane-associated metal-ion transporter that is essential for ethylene signalling (Alonso *et al.*, 1999).

We here demonstrated that SA and JA both act to confer thermotolerance; however, in our experimental approach, ET acts to increase HS-associated cell death.

## Materials and Methods

All chemicals were obtained from Sigma -Aldrich Company Ltd (Gillingham, UK) unless stated otherwise.

### Plant materials and growth conditions

*Arabidopsis thaliana* (L.) Heynh. ecotype Columbia (Col-0) was used as the wild-type control. The mutant lines, *cpr5-1*, *npr1-1*, *jar1-1*, *ein2-1*, *cpr5-1 npr1-1* and *cpr5-1 jar1-1* (from Xinnian Dong, Duke University, Durham, NC, USA), *HSP17.3B-GUS* (Clarke *et al.*, 2004), *coi1-1* (from Prof. John Turner, University of East Anglia, Norwich, UK) and 35S-NahG line 10, from Scott Uknes (Ciba-Geigy Corp, Greensboro, NC, USA; currently Cropsolution Inc., Morrisville, NC, USA), were in this ecotype background. The *PDF1.2-GUS* line (from Willem Broekaert, Katholieke Universiteit Leuven, Belgium; Manners *et al.*, 1998) was in the C24 ecotype background. The *opr3* line (from John Browse and Annick Stintzi, Pullman, Washington State University) was in the WS-0 background.

*Arabidopsis* plants were grown in 9-cm Petri dishes on Murashige and Skoog (MS; Duchefa, Haarlem, the Netherlands) medium (0.8% (w : v) plant agar, 3.0% (w : v) sucrose, pH 5.8) at 22°C with a 16-h photoperiod (65  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ). Stratified seeds were surface-sterilized in 20% (v : v) bleach for 5 min, rinsed twice in sterile H<sub>2</sub>O (Millipore Milli-Q water purification system; Millipore Australia Pty. Ltd, North Ryde, Australia) before plating onto MS agar plates and grown for 3 wk to principal growth stage 1.08 (Boyes *et al.*, 2001).

### Heat treatment of plants

All heat treatments were carried out in the dark to ensure that cell death was a result of increased temperature and not photo-oxidative stress. Unless otherwise specified, Petri dishes containing 3-wk-old plants on MS medium were heated at the appropriate temperature and specified time in a Sanyo MLR-350 environmental test chamber, without light. When electrolyte leakage was measured during the heat treatment, the aerial parts of 3-wk-old plants were placed individually in glass scintillation vials containing 5 ml H<sub>2</sub>O and placed in the environmental chamber without light (< 1  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) at the appropriate temperature.

For the acquired thermotolerance assay, Petri dishes containing 3-wk-old plants grown on MS medium were exposed to a 2 h 45°C HS with (acclimatized) or without (nonaccli-

matized) a pre-treatment. The pretreatment consisted of 90 min at 38°C followed by a 2-h recovery period at 22°C before the HS. Following the 45°C HS, the plants were returned to 22°C.

The hypocotyl elongation assay was carried out as described by Queitsch *et al.* (2000). Stratified seeds were surface-sterilized and plated in rows on MS medium. Plates were covered with foil and placed in a vertical position at 22°C for 2.5 d. The plates were then either kept at 22°C or heat shocked at 45°C with (acclimatized) or without (nonacclimatized) a 90-min pretreatment at 38°C. Hypocotyl elongation was examined after an additional 2.5 d.

### Electrolyte leakage measurements

Heat-induced changes in cell death were quantified by measuring ion leakage with a Horiba Twin Cond B-173 conductivity meter (HORIBA Ltd, Kyoto, Japan). For analysis of electrolyte leakage following a 16-h heat treatment, aerial parts of 3-wk-old plants were placed in glass scintillation vials containing 5 ml H<sub>2</sub>O and were incubated at 22°C in the light (65 μmol m<sup>-2</sup> s<sup>-1</sup>), unless otherwise stated. The conductance of the H<sub>2</sub>O was measured at intervals and calculated per mg fresh weight. Boiling plants of each genotype to determine total ion content established that this did not differ among the plant lines used in this study (data not shown).

### Chemical treatment of plants

For experiments where MeJA was applied to the medium for electrolyte leakage analysis, 3 ml at the appropriate concentration in H<sub>2</sub>O was filter sterilized and pipetted around the plants 24 h before heat treatment. For gene expression analysis, plants were sprayed with either 0 or 5 μM MeJA. The aerial part of the plant was harvested after 30 min, surface moisture gently removed and immediately frozen in liquid N and stored at -80°C.

### Histochemical GUS staining

Plants were placed into staining solution containing 1 mM 5-bromo-4-chloro-3-indolyl glucuronide (first dissolved in dimethyl formamide at 1 mg ml<sup>-1</sup>) in 50 mM NaPO<sub>4</sub> pH 7.0 and 0.1% Triton X-100. The samples were then placed under vacuum twice for 2 min each time to infiltrate the samples and then incubated at 37°C overnight. The chlorophyll was removed by sequential changes in 30, 75 and 95% ethanol. For histochemical staining of heat-induced GUS activity, plants were returned to 22°C following the heat treatment for 24 h before staining.

### RNA extraction

The aerial part of the plants were harvested following the appropriate treatment and immediately frozen in liquid

nitrogen. Total RNA was prepared from samples kept at -80°C using Plant RNA Purification reagent (Invitrogen, Carlsbad, CA, USA) as per manufactures instructions and quantified using a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). The integrity of RNA samples was also assessed by agarose gel electrophoresis.

### Northern blot analysis

Total RNA was separated on a denaturing formaldehyde agarose gel, and transferred to Hybond N membrane (Amersham International Ltd, GE Healthcare Life Sciences, Little Chalfont, UK) as in Draper *et al.* (1998). Hybridization was carried out as in Warner *et al.* (1992) and the cDNA probe was labelled using the Invitrogen Random Primers DNA Labelling System. The *AtHSP17.6* clone GenBank No. X16076 (stock number CD3-5) was obtained from the *Arabidopsis* Biological Resource Center. The *PDF1.2* cDNA was obtained by polymerase chain reaction (PCR) using the primers 5'-CACCCCTTA-TCTTCGCTGCTC-3' and 5'-GTTGCATGATCCATG-TTTGG-3'.

### cDNA synthesis

To eliminate residual genomic DNA, the RNA was treated with RNase free DNase 1 (Roche Diagnostics, Mannheim, Germany). First-strand cDNA was synthesized by reverse transcribing 1 μg of total RNA using Transcriptor Reverse Transcriptase (Roche Diagnostics, Mannheim, Germany) and 3.2 μg of random hexamer primers (Roche Diagnostics) in a 20-μl reaction as per manufacturers instructions. A two-step incubation was performed to allow efficient annealing of the primers (10 min at 25°C, followed by 30 min at 55°C). Following heat inactivation (85°C for 5 min) of the reverse transcriptase, cDNA samples were stored at -20°C.

### Primer designing and real-time PCR analysis

To facilitate the real-time PCR analysis of all the investigated genes under same reaction conditions, primers were designed using OligoPerfect Designer (Invitrogen) under default parameters. The primers were designed spanning an intron to detect any genomic DNA contamination. Real-time PCR was performed on the Roche LightCycler 2.0 using LightCycler Faststart DNA Master<sup>plus</sup> Sybrgreen1 (Roche Diagnostics) as per manufactures instructions. Each 10-μl reaction contained 1 μl of a 1 : 10 dilution of cDNA and 250 nM of each gene-specific primer. *Actin 2* (At3g18780) and the cap-binding protein (*AtCBP20*; At5g44200) were used as the reference gene internal controls for comparison of the target gene transcripts amplified per cDNA sample. The following primers were used for reference gene amplification: *Actin 2*, 5'-CGCTCTTTC-TTTCGAAGCTCAT-3' and 5'-TCCTGCAAATCCAG-CCTTC-3'; *AtCBP20*, 5'-ATCGCTATGGCCGCGATA-3'

and 5'-AGTCACCGCTCTCACGGAAT-3'. The following primers were used for target gene amplification: *PDF1.2* (At5g44420), 5'-CACCTTATCTTCGCTGCTC-3' and 5'-GTTGCATGATCCATGTTTGG-3'; *THI2.1* (At1g72260), 5'-TTCCAAGGGAAGGTGTATGC-3' and 5'-AGAGGT-TTCATGGCACCACA-3'; *VSP1* (At5g24780), 5'-TTTT-ACGCCAAAGGACTTGC-3' and 5'-AATCCCGAG-TTCCAAGAGGT3'. LightCycler Software 4 (Roche Diagnostics) relative quantification analysis module was used to determine the relative fold expression changes between treatments. This software compares the difference between the amplification of target genes to that obtained for the reference genes for each cDNA treatment sample and normalizes to a value of 1 based on the treatment used as the calibrator (0 min heat stressed or 0  $\mu\text{M}$  MeJA treated plants) and takes into account the efficiency of the primer pair. Standard curves were obtained for each primer pair and the following efficiencies and errors, respectively, were obtained; *Actin 2* (1.96; 0.007), *CAB2* (1.97; 0.009), *PDF1.2* (1.84; 0.002), *THI2.1* (1.88; 0.006) and *VSP1* (1.88; 0.002). All the PCRs were performed under the following conditions: 10 min at 95°C followed by 45 cycles of 5 s at 95°C, 5 s at 60°C and 8 s at 72°C in LightCycler capillaries (Roche Diagnostics). The specificity of amplicons was verified by melting curve analysis (65–95°C) after 45 cycles and agarose gel electrophoresis. At least three biological replicates for each sample were used for real-time PCR analysis and at least two technical replicates were analysed for each biological replicate.

#### Measurements of JA, 12-oxo-phytodienoic acid (OPDA) and JA-Ile levels

The plant material was extracted and analysed on a Micromass LCT ESI-MS (Waters Ltd, Manchester, UK) as described in Clarke *et al.* (2004). The JA levels were calculated from the recovery of JA against an added  $d_6$ -SA standard as described in Allwood *et al.* (2006). In addition, JA, OPDA and JA-Ile were extracted and quantified as described elsewhere (Hause *et al.*, 2000; Miersch *et al.*, 2008) using ( $^2\text{H}_6$ )JA, ( $^2\text{H}_5$ )OPDA and JA-( $^2\text{H}_3$ )Leu as standards and a Polaris Q instrument (Thermo-Finnigan; Thermo Fisher Scientific Inc, Waltham, MA, USA) for gas chromatography–mass spectrometry (GC-MS) analysis.

#### Photoacoustic measurements of ET production

Ethylene production was measured using a sensitive laser-based ET detector (type ETD-300; Sensor Sense BV, Nijmegen, the Netherlands) in combination with a gas handling system. The ETD-300 is a state-of-the art ethylene detector based on laser photoacoustic spectroscopy (Harren *et al.*, 2000) that is able to detect on-line *c.* 300 pptv (parts-per-trillion volume,  $1:10^{12}$ ) of ethylene within a 5-s time scale. The gas handling was performed by a valve control box (type VC-6, Sensor Sense

B.V., Nijmegen, the Netherlands), designed for measuring up to six sampling cuvettes per experiment. The valve control box allowed automated sampling of ethylene production at a flow rate of 3 l h<sup>-1</sup> and its transport to the ETD-300 alternately, in succession for 15 min for each cuvette. In this study, four glass cuvettes were used per experiment. One contained a Petri dish with agar and the other three cuvettes contained a Petri dish with 12 3-wk-old *Arabidopsis* seedlings grown on MS medium supplemented with 3% sucrose. Two cuvettes were incubated in a Sanyo MLR-350 environmental test chamber, without light; gas-lines were fed directly from the chamber through the VC-6 to the ETD-300 allowing prolonged monitoring of ethylene production without opening the chamber. The third cuvette was maintained at 22°C and wrapped in aluminium foil to maintain plants in the dark. For a better overview the ethylene emission rate is displayed every 1 h. Each experiment was repeated giving similar results.


## Results

### NPR1- and JAR1-dependent pathways are required for the enhanced basal thermotolerant phenotype observed for *cpr5-1*

To examine the contribution of the SA and JA pathways in the elevated thermotolerance seen with *cpr5-1* electrolyte leakage was measured in the double mutants *cpr5-1 npr1-1* and *cpr5-1 jar1-1* and the corresponding single mutants. Electrolyte leakage represents a rapid, sensitive and quantitative method to assess stress on plant cells. We have previously used electrolyte leakage to assess basal thermotolerance in the SA-signalling mutants *npr1-1* and *cpr5-1* (Clarke *et al.*, 2004). In agreement with Clarke *et al.* (2004), the *cpr5-1* mutant exhibited reduced electrolyte leakage compared with Col-0 during heat stress and the *npr1-1* mutant had an increase in electrolyte leakage (Fig. 1a). However, a similar level of electrolyte leakage was observed for *cpr5-1 npr1-1* and Col-0 suggesting that the enhanced basal thermotolerant phenotype observed for *cpr5-1* is dependent on NPR1. A similar result was shown following heat stress where the electrolyte leakage levels during heat stress were restored to wild-type levels in the *cpr5-1 npr1-1* mutant compared with the reduction detected for *cpr5-1* (Fig. 1b).

Wild-type levels of electrolyte leakage were detected for *jar1-1* both during heat stress and subsequent recovery (Fig. 1c,d). Although this may suggest that JAR1 is not involved in thermotolerance, electrolyte leakage levels were restored to wild-type levels in the *cpr5-1 jar1-1* double mutant. Significantly less electrolyte leakage was detected for *cpr5-1* compared with that observed for the double mutant *cpr5-1 jar1-1* during heat stress after 5 h ( $P < 0.054$ ) and 7 h ( $P < 0.051$ ; Fig. 1c, and 3 h ( $P < 0.043$ ) and 5 h ( $P < 0.049$ ) post heat stress (Fig. 1d). This could indicate that the JA pathway is required for the enhanced thermotolerance of *cpr5-1*.

**Fig. 1** *JAR1* and *NPR1* are required for the enhanced thermotolerant phenotype of *cpr5-1*. (a,c) Electrolyte leakage in *Arabidopsis thaliana* wild-type Col-0 and the mutants *cpr5-1*, *npr1-1*, *jar1-1* and the double mutants *cpr5-1 npr1-1* and *cpr5-1 jar1-1* during heat treatment. The aerial parts of the plants were incubated in H<sub>2</sub>O at 38°C and the conductivity of the solution was monitored. Data are means ( $\pm$  SE) of six experiments each with five plants per treatment ( $n = 6$ ). (b,d) Electrolyte leakage in wild-type Col-0 and the mutants *cpr5-1*, *npr1-1*, *jar1-1* and the double mutants *cpr5-1 npr1-1* and *cpr5-1 jar1-1* during recovery from heat treatment. Following heat treatment (16 h at 38°C) the aerial parts of the plants were placed in H<sub>2</sub>O at 22°C and the conductivity of the solution was monitored. Data are means ( $\pm$  SE) of six experiments each with five plants per treatment ( $n = 6$ ).



When acquired thermotolerance was assessed in these genotypes, all exhibited this effect to levels equivalent to wild type controls (see the Supporting Information, Fig. S1).

### Heat stress induces the JA signal transduction pathway

Given the equivocal data obtained with the *cpr5-1 jar1-1* double mutant we sought to further investigate the possible role of jasmonates in thermotolerance. Thus, we investigated thermotolerance in *coi1-1*, which plays a more central role in the signalling mechanism (Staswick, 2008). A clear increase in electrolyte leakage was observed for *coi1-1* compared with that observed for Col-0 both during the 38°C heat treatment and subsequent recovery from a 16-h 38°C heat treatment (Fig. 2a,b). It was also noted that unlike any other mutant genotype, *coi1-1* plants were found to have severe chlorosis beginning from the base of the petioles (Fig. 2c).


We sought further evidence for the involvement of the jasmonates in heat stress in wild-type plants. The JA-inducible genes *PDF1.2* (At5g44420), *THI2.1* (At1g72260) and *VSP1* (At5g24780) were analysed in wild-type (Col-0) plants following a 38°C heat treatment. Heat-induced expression was initially assessed by Northern analysis but while *PDF1.2* transcript accumulation was noted no expression was observed for *THI2.1* or *VSP1* (data not shown). To confirm this result, real-time PCR was employed. A threefold increase in *PDF1.2*

transcript was detected following a 30-min heat treatment at 38°C and this remained relatively constant during the 360-min heat treatment (Fig. 3a). By contrast, transcript levels at only 45% and 37% those of controls were detected with *THI2.1* and *VSP1*, respectively, following a 30-min heat stress. Following 360 min of heat stress both *THI2.1* and *VSP1* transcript levels were approx. 10% those of nonHS plants.


In investigating the reason for this differential response of jasmonate marker genes we examined whether they exhibited differing sensitivities to jasmonate. *PDF1.2* displayed a greater response to exogenous application of 5  $\mu$ M MeJA where a 300-fold increase over controls was detected, compared with a 3- and 36-fold increase for *THI2.1* and *VSP1* transcripts, respectively (Fig. 3b).

Plants transformed with *GUS* driven by the *PDF1.2* promoter were employed to investigate the spatial patterns of heat-induced *PDF1.2* expression. Although basal level of *GUS* expression was detected throughout the plant, an increase in *GUS* expression was detected following 2 h of heat stress (Fig. 3c). The intensity of the heat-induced *GUS* expression increased following 6 h and 12 h of heat stress and appeared to be localized to leaf tissue with little upregulation in root or stem tissue.

Further support for a protective role of jasmonate during heat stress was shown when 3-wk-old Col-0 plants were subjected to a 38°C heat treatment following exogenous application




**Fig. 2** *coi1-1* is more susceptible to heat stress. (a) Electrolyte leakage during heat treatment. The aerial parts of *Arabidopsis thaliana* Col-0 (squares) and *coi1-1* (triangles) were incubated in H<sub>2</sub>O at 38°C and the conductivity of the solution was monitored. Data are means (± SE) of four experiments each with five plants per treatment ( $n = 4$ ). (b) Electrolyte leakage during recovery from heat treatment. Following heat treatment (16 h; 38°C) the aerial parts of Col-0 (squares) and *coi1-1* (triangles) were placed in H<sub>2</sub>O at 22°C and the conductivity of the solution was monitored. Data are means (± SE) of five experiments each with five plants per treatment ( $n = 5$ ). (c) Col-0 (left) and *coi1-1* (right) 3-wk-old plants on Murashige and Skoog (MS) medium were exposed to 38°C for 16 h, returned to 22°C and photographed 4 d later.



**Fig. 3** Heat induces *PDF1.2* expression. (a) *PDF1.2* (closed bars), *THI2.1* (open bars) and *VSP1* (tinted bars) transcript accumulation in *Arabidopsis thaliana* Col-0 during heat stress. Real-time polymerase chain reaction (PCR) was performed on cDNA synthesised from total RNA extracted from the aerial parts of 3-wk-old Col-0 exposed to a 38°C heat treatment for the specified time. Transcript expression was normalized to the amount detected for nonheat stressed tissue. Data are means (± SE) of at least five experiments ( $n \geq 5$ ). (b) Induction of *PDF1.2*, *THI2.1* and *VSP1* transcripts in Col-0 following methyl jasmonate (MeJA) treatment. Real-time PCR performed on cDNA synthesized from total RNA extracted from the aerial parts of 3-wk-old Col-0 sprayed with 5 μM methyl jasmonate (MeJA). Transcript expression was normalized to the amount detected for 0 μM MeJA treated tissue. Data are means (± SE) of at least five experiments ( $n \geq 5$ ). (c) *PDF1.2-GUS* expression during heat stress: 3-wk-old *PDF1.2-GUS* plants were incubated at 38°C for 0, 2, 6 and 12 h, and returned to 22°C for 24 h before staining.

of MeJA 24 h before heat stress. Electrolyte leakage measurements indicated that treatment with 1 μM MeJA did not provide any protection from heat stress but 5 μM MeJA significantly ( $P < 0.013$ ) reduced electrolyte leakage compared with the

control plants (Fig. 4a). However, further increases in MeJA concentration had the reverse effect, suggesting that MeJA protection from heat stress is concentration specific. To confirm that exogenous application of MeJA via the media was inducing



**Fig. 4** Methyl jasmonate (MeJA) protects *Arabidopsis thaliana* Col-0 from heat stress. (a) H<sub>2</sub>O (0 μM MeJA), 1 μM MeJA, 5 μM MeJA, 10 μM MeJA and 30 μM MeJA was applied to the Murashige and Skoog (MS) media 24 h before heat treatment (16 h for 38°C). Following heat treatment the aerial parts were placed in H<sub>2</sub>O at 22°C and the conductivity of the solution was monitored. Data are means (± SE) of four experiments each with five plants per treatment ( $n = 4$ ). (b) Northern analysis of *PDF1.2* and *HSP17.6* transcripts in 3-wk-old Col-0 seedlings exposed to the various concentrations (μM) of MeJA indicated by supplementing the culture media. Northern blots of separated RNA (10 μg) were hybridized with <sup>32</sup>P-labelled *PDF1.2* or *HSP17.6* cDNA. RNA (2 μg) from Col-0 seedlings exposed to a heat treatment (38°C for 60 min) was included as a positive control for *HSP17.6* transcripts. The ethidium bromide-stained gel is shown below.

the JA pathway, the expression of *PDF1.2* was assessed by Northern analysis. This indicated that exogenous application of MeJA induced *PDF1.2*.

When we assessed the expression of *HSPs* that are well-established markers for thermotolerance (Clarke *et al.*, 2004), MeJA did not induce expression of *HSP17.6* (At5g12020; Fig. 4b); neither was *HSP101* (At1g74310) detected (data not shown). No GUS expression was detected in *HSP17.3*-GUS plants following exogenous application whereby MeJA was either pipetted onto the media or the plant was submerged in a solution of MeJA (data not shown). Examination of publicly available array data using GENEVESTIGATOR (Zimmermann

*et al.*, 2004) also did not indicate that MeJA induced any class of HSP (Fig. S2). Hence, we can find no evidence that MeJA-conferred thermotolerance is associated with HSP gene expression.


Next, we analysed the endogenous free JA levels in Col-0 during heat stress (Fig. 5a). The JA levels were significantly ( $P \leq 0.001$ ) increased over levels at  $t = 0$  at 1 h of 38°C heat treatment and continued to rise afterwards. To indicate the contribution of *de novo* JA biosynthesis to thermotolerance, heat tolerance in the OPDA reductase 3 mutant, *opr3* was examined. During heating no significant difference was observed between Col-0 and *opr3* (Fig. 5b) but during the recovery phase, *opr3* was found to be compromised, exhibiting greater electrolyte leakage (Fig. 5c).

To relate these *opr3* responses with jasmonates; levels of JA, OPDA and JA-Ile were determined (Fig. 5d–f). Significant accumulation of all jasmonates was observed in response to heating to 38°C by 8 h. In the case of JA and OPDA levels were significantly reduced at 8 h following heating in *opr3*. The JA-Ile levels in *opr3* were significantly reduced before and following heating to 38°C until 12 h (Fig. 5f).

#### Heat stress initiates ethylene production

The expression of *PDF1.2* is influenced by both JA and ET (Pré *et al.*, 2008), hence the significant elevation of its expression during HS could indicate a contribution by ET. Larkindale *et al.* (2005) demonstrated that following heating to 45°C for 60 min, the *ein2-1* mutant was compromised in its tolerance to heat. By contrast, with our gentler heating regime dramatically different responses were observed with the *ein2-1* mutant. Electrolyte leakage was decreased in *ein2-1* compared with Col-0 following heat stress (Fig. 6a) but not during heating (Fig. 6b) suggesting that ET contributed to the poststress loss in viability.

To investigate this further, ET production in 2-wk-old agar-grown seedlings was measured by laser photoacoustic detection (Fig. 6c). Since the actual ET concentration levels were far below 1 ppbv (parts-per-billion volume,  $1:10^9$ ) this technique had to be employed instead of the traditional gas chromatography. Incubation of Col-0 at 38°C resulted in a rapid increase in ET production that persisted for at least 24 h. Ethylene production from the agar was negligible. The possible contribution of JA and SA to ET production was assessed using *opr3* and a line harbouring the SA hydroxylase *NahG* transgene that are unable to accumulate SA (Delaney *et al.*, 1994). When *opr3* mutants were incubated at 38°C, ET production was reduced compared with wild-type (*Ws-0*) plants (Fig. 6d). Examination of ET production in *NahG* plants at 38°C was also reduced compared with wild type (Col-0) controls (Fig. 6e). Taking these data together it appears that both JA and SA augment ET production in response to heating. However, ET is likely to be exerting a negative effect on plant cell survival.



**Fig. 5** Levels of jasmonic acid (JA), 12-oxophytodienoic acid (OPDA) and JA-isoleucine conjugate (JA-Ile) rise during heat stress. (a) Endogenous levels of JA in Col-0 plant tissue exposed to 38°C for the indicated time. Data are means ( $\pm$  SE) of at least three replicates. (b) Electrolyte leakage during heat treatment. The aerial parts of the plants were incubated in H<sub>2</sub>O at 38°C and the conductivity of the solution was monitored. Data are means ( $\pm$  SE) of six experiments each with five plants per treatment ( $n = 6$ ). (c) Electrolyte leakage during recovery from heat treatment of Ws-0 (squares) and *opr3* (triangles) mutant. Following heat treatment (16 h for 38°C) the aerial parts of the plants were placed in H<sub>2</sub>O at 22°C and the conductivity of the solution was monitored. Data are means ( $\pm$  SE) of six experiments each with five plants per treatment ( $n = 6$ ). (d–f) Levels of JA, OPDA and JA-Ile in wild type (Ws-0; tinted bars) and *opr3* (open bars) plants. Data are three replicate plants per treatment. Error bars are  $\pm$  SE. Significance: \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ ; ns, not significant.


## Discussion

This study developed observations made in our previous study (Clarke *et al.*, 2004) where the *Arabidopsis* mutant *cpr5-1* displayed exceptional thermotolerance. Electrolyte leakage, as a rapidly measured parameter, enabled us to establish that the enhanced thermotolerant phenotype observed for *cpr5-1* *Arabidopsis* plants was compromised in the *cpr5-1 npr1-1* double mutant. This indicated that NPR1 is required for the enhanced basal thermotolerance observed for *cpr5-1*. Studies into induced systemic resistance (ISR) have indicated that NPR1 also acts to integrate defence signalling by SA, JA and ET (Pieterse & van Loon, 2004). We have previously indicated that SA plays a role in basal thermotolerance but the *npr1* data could indicate contributions by JA or ET. Crucially, however, an NPR1-independent pathway must also exist because the electrolyte leakage detected for the double mutant was similar to that observed for Col-0 and not *npr1-1*, which exhibited a higher level of electrolyte leakage.

Clarke *et al.* (2000) established that the *cpr5-1* phenotype in biotic stress responses is partially influenced by the JA-signalling pathway, so we first examined if this could also be contributing to the thermotolerance pathway. To investigate this possibility we employed *jar1-1* and the *cpr5-1 jar1-1* double mutant. The single *jar1-1* mutant was not compromised in

basal thermotolerance suggesting that JAR1 is not required for tolerance during heating. However, the enhanced thermotolerance observed in *cpr5-1* was not seen with the *cpr5-1 jar1-1* double mutant implying a requirement for jasmonates. Based on these somewhat equivocal observations, we tested the role of JA in thermotolerance within wild-type *Arabidopsis*. Electrolyte leakage experiments demonstrated that exogenous application of 5  $\mu$ M MeJA helped to maintain cell viability in heat-stressed plants. However, increasing the concentration to 10  $\mu$ M had no protective affect and a further increase to 30  $\mu$ M resulted in increased membrane damage. Given that this experiment supplemented endogenous JA production, this suggested a threshold for jasmonate-conferred thermotolerance. Further, jasmonate increases occur at the same time as SA production, this could be an example of the reduced viability seen with high concentration SA–JA interactions (Mur *et al.*, 2006). However, when the expression of JA-inducible genes *PDF1.2*, *THI2.1* and *VSP1* during heat stress were monitored a threefold induction of only *PDF1.2* transcript was detected whereas *THI2.1* and *VSP1* transcripts decreased during heat stress. One likely explanation is that both *THI2.1* and *VSP1* are less sensitive to MeJA, as only a 3- and 36-fold increase in transcript was found, respectively, following MeJA treatment compared with a 300-fold increase for *PDF1.2*. The endogenous level of JA detected during heat stress may therefore be





**Fig. 6** Analysis of heat-treated *ein2-1* and ethylene production in *Arabidopsis thaliana* wild type, *opr3* and *NahG* *Arabidopsis* plants during heating to 38°C. (a) Electrolyte leakage during recovery from heat treatment. Following heat treatment (16 h for 38°C) the aerial parts of the plants were placed in H<sub>2</sub>O at 22°C and the conductivity of the solution was monitored. Data are means ( $\pm$  SE) of six experiments each with five plants per treatment ( $n = 6$ ). (b) Electrolyte leakage during heat treatment. The aerial parts of the plants were incubated in H<sub>2</sub>O at 38°C and the conductivity of the solution was monitored. Data are means ( $\pm$  SE) of six experiments each with five plants per treatment ( $n = 6$ ). (c–e) C<sub>2</sub>H<sub>4</sub> production from 12 2-wk-old *Arabidopsis* seedlings, grown on Murashige and Skoog (MS) medium + 3% sucrose following heating to 38°C and measured using laser photoacoustic detection. (c) C<sub>2</sub>H<sub>4</sub> produced from Col-0 incubated at 38°C from two separate experiments (diamonds, triangles); C<sub>2</sub>H<sub>4</sub> from Col-0 incubated at 22°C (squares) and from Petri-dishes with MS salts heated to 38°C (circles). (d) C<sub>2</sub>H<sub>4</sub> produced from *Ws-0* incubated at 38°C (closed diamonds); C<sub>2</sub>H<sub>4</sub> from *opr3* incubated at 38°C from two separate experiment (open diamonds, squares) and from Petri-dishes with MS salts heated to 38°C (circles). (e) C<sub>2</sub>H<sub>4</sub> produced from Col-0 incubated at 38°C (diamonds); C<sub>2</sub>H<sub>4</sub> from *NahG* incubated at 38°C from two separate experiment (closed circles, triangles) and from Petri dishes with MS salts heated to 38°C (open circles). Seedlings were maintained in the dark for all measurements.

inadequate to induce *THI2.1* and *VSP1* transcription. Contrasting expression of *PDF1.2* compared with both *THI2.1* and *VSP1* genes has also been seen in response to JA via the transcription factor AtMYC2 which upregulated wound-response genes *VSP* and *THI2* but repressed *PDF1.2* (Lorenzo *et al.*, 2004). However, microarray analysis of MeJA-treated *Arabidopsis* seedlings grown on MS-agar failed to exhibit significant *PDF1.2* expression. This was in contrast to both *VSP1* and *THI2.1* which were readily induced by MeJA (Devoto *et al.*, 2005). Instead, high-level *PDF1.2* expression was dependent on both jasmonates and ET (Penninckx *et al.*, 1998). Hence, the rapid induction of *PDF1.2* on heating

(Fig. 3a) may be a reflection of both jasmonate (Fig. 5) and ET production (Fig. 6).

Although, exogenous application of MeJA protects membranes from heat stress, categorical proof of a role for jasmonates in thermotolerance required a JA mutant to demonstrate reduced heat-resistance. Unlike *jar1-1*, *coi-1* is defective in many JA-dependent responses (Xie *et al.*, 1998, Farmer *et al.*, 2003) and, crucially, we found *coi-1* to be thermosensitive to heat stress. The *COI1-1* gene encodes an F-box protein, part of the SCF<sup>COI</sup> (Skip-cullin-F-box) ubiquitin ligase complex, providing the first indication that ubiquitin-mediated protein degradation is involved in JA signalling (Xie *et al.*, 1998; Xu

*et al.*, 2002; Devoto *et al.*, 2003; Feng *et al.*, 2003). Temperature stress, both high and low, has also been shown to induce transcript of other E3 ubiquitin ligases (Yan *et al.*, 2003). In an attempt to generate plants with increased stress tolerance, Yan *et al.* (2003) overexpressed the *AtCHIP* gene (encoding a protein with a U-box domain and E3 ubiquitin ligase activity) in *Arabidopsis*. However, this rendered the plant more sensitive to both low- and high-temperature treatment, with an increase in electrolyte leakage observed (Yan *et al.*, 2003); this suggests that expression of this E3 ubiquitin ligase at least, is under tight regulation and any disturbances lead to serious consequences in temperature stress tolerance. This is highlighted by our finding that the *coi1-1* plants, impaired in the ubiquitin-mediated degradation pathway, also show increased heat-induced membrane damage.

A significant finding of our study was the discrete roles for particular jasmonates in thermotolerance. MeJA/JA could protect plants during both heating and recovery phases (Fig. 2a,b), but not confer acquired thermotolerance. However, only recovery following stress (Fig. 5b) was compromised with the *opr3* (OPDA reductase) mutant but not leakage during heating (Fig. 5a). This suggests the OPDA was sufficient for basal thermotolerance but downstream jasmonates are required for the recovery phase. Significantly, in a comparison of OPDA versus MeJA induced gene expression, only OPDA proved to be sufficient to induce a range of heat shock proteins (Taki *et al.*, 2005).

An additional signalling interaction within *cpr5-1* was that between SA and ET. In contrast to the findings of Larkindale *et al.* (2005), we found that *ein2-1* plants were less susceptible to heat stress. The discrepancy observed may be caused by the difference in heat treatment and age of the plants; Larkindale *et al.* (2005) subjected 7-d-old seedlings to a 60-min 45°C heat treatment to assess basal thermotolerance compared with our milder experimental condition of 38°C heat treatment for 16°h using 3-wk-old plants. One interpretation based on evidence from the literature is that ET is a response to stress and an indicator of cell death. Ethylene has been linked to the initiation of cell death associated with, for example, ozone (Vahala *et al.*, 2003), disease (Cohn & Martin, 2005) and in roots (He *et al.*, 1996). Further, this matches our own observations of reduced cell death in *ein2-1*. Moreover, with NahG and *opr3*, where heat tolerance is reduced and cell death increased (Clarke *et al.*, 2004; Fig. 5c), ET levels are lower (Fig. 6). Hence, the ET may simply reflect the levels of cell death at 38°C but given the data of Larkindale *et al.* (2005) ET may act to confer thermotolerance at higher temperatures. This latter effect may reflect differential ET interactions with SA and JA. This important aspect requires further exploration. It is interesting to note that *ein2-1* and *coi1-1* mutations appear to have opposite effects on thermotolerance in 3-wk-old plants. Jasmonic acid and ET can either cooperate or act as antagonists in the regulation of different stress responses. In the case of pathogen attack, both hormones appear to synergize

in the activation of defence gene expression (Xu *et al.*, 1994; Penninckx *et al.*, 1998; Lorenzo *et al.*, 2003). However, an antagonistic interaction between JA and ET has been described in response to ozone exposure where JA protects tissues from ozone stress and ET enhances ozone-induced cell death (Overmyer *et al.*, 2000; Tamaoki *et al.*, 2003; Tuominen *et al.*, 2004). In addition the *jar1-1 Arabidopsis* mutant was found to be ozone sensitive in contrast to *ein2-1*, which is ozone tolerant (Tuominen *et al.*, 2004). Such ET–JA antagonism may effect at the transcriptional level as MYC2-mediated induction of *VSP1* is suppressed by ET (Dombrecht *et al.*, 2007). This contrasts with the positive contribution made by JA to *PDF1.2* expression acting through the ERF1 and ORA59 transcription factors which were first characterized as mediating ET effects (Pré *et al.*, 2008). An important task for future work is to establish the mechanism(s) for JA-conferred thermotolerance, given that HSP apparently play no role (Fig. 4b, Fig. S2). Examination of transcriptome data for heat and MeJA responsive genes via GENEVESTIGATOR (Zimmermann *et al.*, 2004) indicated a preponderance of genes associated with protein translation and photosynthetic electron transport, which could represent features associated with cellular recovery following HS (Fig. S3).

Through a network of signalling pathways plants are able to tolerate both biotic and abiotic stresses where selection of the specific response that is appropriate for each stress stimulus seems to be determined by the type of interaction established between the signalling pathways. Our studies (this and Clarke *et al.*, 2004) and those of Larkindale & Knight (2002) and Larkindale *et al.* (2005) shed some light on crosstalk between the hormonal, redox and calcium-activated signalling pathways which play a role in thermotolerance to ensure a rapid response which is crucial for survival against temperature stress.

## Acknowledgements

We are extremely grateful for the materials donated by Xinnian Dong (Duke University, NC, USA), Willem Broekaert (CropDesign NV, Belgium) and John Browse as well as Annick Stintzi (Washington State University, Pullman, WA, USA). We thank Jim Heald (Aberystwyth University, UK) for operation of the LC-MS and William Allwood (Manchester University, UK) for valuable assistance with JA extraction and analysis. We also appreciate the constructive comments made by the anonymous reviewers. This work was supported by a NZ Science and Technology Post-Doctoral Fellowship to S.M.C. and the EU-FP6-Infrastructures-5 program, project FP6-026183 'Life Science Trace Gas Facility'.

## References

- Allwood JW, Ellis DI, Heald JK, Goodacre R, Mur LAJ. 2006. Metabolomic approaches reveal that phosphatidic and phosphatidyl

- glycerol phospholipids are major discriminatory metabolites in responses by *Brachypodium distachyon* to challenge by *Magnaporthe grisea*. *Plant Journal* 46: 351–368.
- Alonso JM, Hirayama T, Roman G, Nourizadeh S, Ecker JR. 1999. EIN2-1, a bifunctional transducer of ethylene and stress responses in *Arabidopsis*. *Science* 284: 2148–2152.
- Balbi V, Devoto A. 2008. Jasmonate signalling network in *Arabidopsis thaliana*: crucial regulatory nodes and new physiological scenarios. *New Phytologist* 177: 301–318.
- Berger S, Bell E, Mullet JE. 1996. Two methyl jasmonate-insensitive mutants show altered expression of *AtVsp* in response to methyl jasmonate and wounding. *Plant Physiology* 111: 525–531.
- Boston RS, Viitanen PV, Vierling E. 1996. Molecular chaperones and protein folding in plants. *Plant Molecular Biology* 32: 191–222.
- Bowling SA, Clarke JD, Liu YD, Klessig DF, Dong X. 1997. The *cpr5-1* mutant of *Arabidopsis* expresses both NPR1-dependent and NPR1-independent resistance. *Plant Cell* 9: 1573–1584.
- Boyes DC, Zayed AM, Ascenzi R, McCaskill AJ, Hoffman NE, Davis KR, Gortlach J. 2001. Growth stage-based phenotypic analysis of *Arabidopsis*: a model for high throughput functional genomics in plants. *The Plant Cell* 13: 1499–1510.
- Chini A, Fonseca S, Fernández G, Adie B, Chico JM, Lorenzo O, García-Casado G, López-Vidriero I, Lozano FM, Ponce MR *et al.* 2007. The JAZ family of repressors is the missing link in jasmonate signalling. *Nature* 448: 666–671.
- Clarke JD, Liu YD, Klessig DF, Dong XN. 1998. Uncoupling PR gene expression from NPR1 and bacterial resistance: characterization of the dominant *Arabidopsis cpr6-1* mutant. *Plant Cell* 10: 557–569.
- Clarke JD, Volko SM, Ledford H, Ausubel FM, Dong XN. 2000. Roles of salicylic acid, jasmonic acid, and ethylene in cpr-induced resistance in *Arabidopsis*. *The Plant Cell* 12: 2175–2190.
- Clarke SM, Mur LAJ, Wood JE, Scott IM. 2004. Salicylic acid dependent signalling promotes basal thermotolerance but is not essential for acquired thermotolerance in *Arabidopsis thaliana*. *Plant Journal* 38: 432–447.
- Cohn JR, Martin GB. 2005. *Pseudomonas syringae* pv. *tomato* type III effectors *AvrPto* and *AvrPtoB* promote ethylene-dependent cell death in tomato. *Plant Journal* 44: 139–154.
- Dat JF, Foyer CH, Scott IM. 1998b. Changes in salicylic acid and antioxidants during induced thermotolerance in mustard seedlings. *Plant Physiology* 118: 1455–1461.
- Dat JF, Lopez-Delgado H, Foyer CH, Scott IM. 1998a. Parallel changes in H<sub>2</sub>O<sub>2</sub> and catalase during thermotolerance induced by salicylic acid or heat acclimation in mustard seedlings. *Plant Physiology* 116: 1351–1357.
- Delaney TP, Uknes S, Vernooij B, Friedrich L, Weymann K, Negrotto D, Gaffney T, Gut-Rella M, Kessmann H, Ward E *et al.* 1994. A central role of salicylic acid in plant disease resistance. *Science* 266: 1247–1250.
- Devadas SK, Enyedi A, Raina R. 2002. The *Arabidopsis hrll* mutation reveals novel overlapping roles for salicylic acid, jasmonic acid and ethylene signalling in cell death and defence against pathogens. *Plant Journal* 30: 467–480.
- Devoto A, Ellis C, Magusin A, Chang H-S, Chilcott C, Zhu T, Turner JG. 2005. Expression profiling reveals *COI1* to be a key regulator of genes involved in wound- and methyl jasmonate-induced secondary metabolism, defence, and hormone interactions. *Plant Molecular Biology* 58: 497–513.
- Devoto A, Muskett PR, Shirasu K. 2003. Role of ubiquitination in the regulation of plant defence against pathogens. *Current Opinions in Plant Biology* 6: 307–311.
- Doares SH, Narvaez-Vasquez J, Conconi A, Ryan CA. 1995. Salicylic-acid inhibits synthesis of proteinase-inhibitors in tomato leaves induced by systemin and jasmonic acid. *Plant Physiology* 108: 1741–1746.
- Dombrecht B, Xue GP, Sprague SJ, Kirkegaard JA, Ross JJ, Reid JB, Fitt GP, Sewelam N, Schenk PM, Manners JM *et al.* 2007. MYC2 differentially modulates diverse jasmonate-dependent functions in *Arabidopsis*. *The Plant Cell* 19: 2225–2245.
- Dong X. 2001. Genetic dissection of systemic acquired resistance. *Current Opinions in Plant Biology* 4: 309–314.
- Draper J, Scott R, Armitage P, Walden R. 1998. *Plant genetic transformation and gene expression: a laboratory manual*. Oxford, UK: Blackwell Scientific Publications.
- Farmer EE, Almeras E, Krishnamurthy V. 2003. Jasmonates and related oxylipins in plant responses to pathogenesis and herbivory. *Current Opinions in Plant Biology* 6: 372–378.
- Feng S, Ma L, Wang X, Xie D, Dinesh-Kumar SP, Wei N, Deng XW. 2003. The COP9 signalosome interacts physically with SCF<sup>COI1-1</sup> and modulates jasmonate responses. *Plant Cell* 15: 1083–1094.
- Foyer CH, Lopez-Delgado H, Dat JF, Scott IM. 1997. Hydrogen peroxide and glutathione-associated mechanisms of acclimatory stress tolerance and signaling. *Physiologia Plantarum* 100: 241–254.
- Harren FJM, Cotti G, Oomens J, te Lintel Hekkert S. 2000. Photoacoustic spectroscopy in trace gas monitoring. In: Meyers RA, ed, *Encyclopedia of analytical chemistry*. Chichester, UK: John Wiley, 2203–2226.
- Hause B, Stenzel I, Miersch O, Maucher H, Kramell R, Ziegler J, Wasternack C. 2000. Tissue-specific oxylipin signature of tomato flowers – allene oxide cyclase is highly expressed in distinct flower organs and vascular bundles. *Plant Journal* 24: 113–126.
- He CJ, Morgan PW, Drew MC. 1996. Transduction of an ethylene signal is required for cell death and lysis in the root cortex of maize during aerenchyma formation induced by hypoxia. *Plant Physiology* 112: 463–472.
- Hilpert B, Bohlmann H, op den Camp RO, Przybyla D, Miersch O, Buchala A, Apel K. 2001. Isolation and characterization of signal transduction mutants of *Arabidopsis thaliana* that constitutively activate the octadecanoid pathway and form necrotic microlesions. *Plant Journal* 26: 435–446.
- Hong SW, Lee U, Vierling E. 2003. *Arabidopsis hot* mutants define multiple functions required for acclimation to high temperatures. *Plant Physiology* 132: 757–767.
- Hong SW, Vierling E. 2000. Mutants of *Arabidopsis thaliana* defective in the acquisition of tolerance to high temperature stress. *Proceedings of the National Academy of Sciences, USA* 97: 4392–4397.
- Howe GA. 2004. Jasmonates as signals in the wound response. *Journal of Plant Growth Regulation* 23: 223–237.
- Kanna M, Tamaoki M, Kubo A, Nakajima N, Rakwal R, Agrawal GK, Tamogami S, Ioki M, Ogawa D, Saji H *et al.* 2003. Isolation of an ozone-sensitive and jasmonate-semi-insensitive *Arabidopsis* mutant (*oji1*). *Plant Cell Physiology* 44: 1301–1310.
- Katsir L, Chung HS, Koo AJK, Howe GA. 2008. Jasmonate signaling: a conserved mechanism of hormone sensing. *Current Opinion in Plant Biology* 11: 1–8.
- Larkindale J, Hall JD, Knight MR, Vierling E. 2005. Heat stress phenotypes of *Arabidopsis* mutants implicate multiple signaling pathways in the acquisition of thermotolerance. *Plant Physiology* 138: 882–897.
- Larkindale J, Knight MR. 2002. Protection against heat stress-induced oxidative damage in *Arabidopsis* involves calcium, abscisic acid, ethylene, and salicylic acid. *Plant Physiology* 128: 682–695.
- Laudert D, Weiler EW. 1998. Allene oxide synthase: a major control point in *Arabidopsis thaliana* octadecanoid signalling. *Plant Journal* 15: 675–684.
- Lorenzo O, Chico JM, Sanchez-Serrano JJ, Solano R. 2004. Jasmonate-insensitive1 encodes a MYC transcription factor essential to discriminate between different jasmonate-regulated defence responses in *Arabidopsis*. *The Plant Cell* 16: 1938–1950.
- Lorenzo O, Piqueras R, Sanchez-Serrano JJ, Solano R. 2003. ETHYLENE RESPONSE FACTOR1 integrates signals from ethylene and jasmonate pathways in plant defense. *The Plant Cell* 15: 165–178.
- Love AJ, Yun BW, Laval V, Loake GJ, Milner JJ. 2005. Cauliflower mosaic virus, a compatible pathogen of *Arabidopsis*, engages three distinct defense-signaling pathways and activates rapid systemic generation of reactive oxygen species. *Plant Physiology* 139: 935–948.

- Manners JM, Penninckx IAM, Vermaere K, Kazan K, Brown RL, Morgan A, Maclean DJ, Curtis MD, Cammue BPA, Broekaert WF. 1998. The promoter of the plant defensin gene PDF1.2 from *Arabidopsis* is systemically activated by fungal pathogens and responds to methyl jasmonate but not to salicylic acid. *Plant Molecular Biology* 38: 1071–1080.
- McConn M, Creelman RA, Bell E, Mullet JE, Browse J. 1997. Jasmonate is essential for insect defense in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* 94: 5473–5477.
- Miersch O, Neumerkel J, Dippe M, Stenzel I, Wasternack C. 2008. Hydroxylated jasmonates are commonly occurring metabolites of jasmonic acid and contribute to a partial switch-off in jasmonate signaling. *New Phytologist* 177: 114–127.
- Mur LAJ, Kenton P, Atzorn R, Miersch O, Wasternack C. 2006. The outcomes of concentration-specific interactions between salicylate and jasmonate signaling include synergy, antagonism, and oxidative stress leading to cell death. *Plant Physiology* 140: 249–262.
- Navarre DA, Wolpert TJ. 1999. Victorin induction of an apoptotic/senescence-like response in oats. *Plant Cell* 11: 237–49.
- Overmyer K, Tuominen H, Kettunen R, Betz C, Langebartels C, Sandermann H Jr, Kangasjärvi J. 2000. Ozone-sensitive *Arabidopsis rcd1* mutant reveals opposite roles for ethylene and jasmonate signaling pathways in regulating superoxide-dependent cell death. *The Plant Cell* 12: 1849–1862.
- Pastori GM, Foyer CH. 2002. Common components, networks, and pathways of cross-tolerance to stress. The central role of 'redox' and abscisic acid-mediated controls. *Plant Physiology* 129: 460–468.
- Pěna-Cortes H, Albrecht T, Prat S, Weiler EW, Willmitzer L. 1993. Aspirin prevents wound-induced gene-expression in tomato leaves by blocking jasmonic acid biosynthesis. *Planta* 191: 123–128.
- Penninckx IAM, Thomma BPHJ, Buchala A, Mettraux J-P, Broekaert WF. 1998. Concomitant activation of jasmonate and ethylene response pathways is required for induction of a plant defensin. *The Plant Cell* 10: 2103–2113.
- Pieterse CM, van Loon LC. 2004. NPR1: the spider in the web of induced resistance signaling pathways. *Current Opinions of Plant Biology* 7: 456–464.
- Pré M, Atallah M, Champion A, De Vos M, Pieterse CMJ, Memelink J. 2008. The AP2/ERF-domain transcription factor ORA59 integrates jasmonic acid and ethylene signals in plant defense. *Plant Physiology* 147: 1347–1357.
- Queitsch C, Hong S-W, Vierling E, Lindquist S. 2000. Hsp101 plays a crucial role in thermotolerance in *Arabidopsis*. *Plant Cell* 12: 479–492.
- Rao MV, Lee H, Creelman RA, Mullet JE, Davis KR. 2000. Jasmonic acid signaling modulates ozone-induced hypersensitive cell death *Plant Cell* 12: 1633–1646.
- Reymond P, Bodenhausen N, van Poecke RM, Krishnamurthy V, Dicke M, Farmer EE. 2004. A conserved transcript pattern in response to a specialist and a generalist herbivore. *Plant Cell* 16: 3132–3147.
- Reymond P, Weber H, Damond M, Farmer EE. 2000. Differential gene expression in response to mechanical wounding and insect feeding in *Arabidopsis*. *Plant Cell* 12: 707–720.
- Spoel SH, Koornneef A, Claessens SMC, Korzelius JP, van Pelt JA, Mueller MJ, Buchala AJ, Mettraux JP, Brown R, Kazan K *et al.* 2003. NPR1 modulates cross-talk between salicylate- and jasmonate-dependent defense pathways through a novel function in the cytosol. *Plant Cell* 15: 760–770.
- Staswick P. 2008. JAZing up jasmonate signaling. *Trends in Plant Science* 13: 66–71.
- Staswick PE, Tiryaki I. 2004. The oxylipin signal jasmonic acid is activated by an enzyme that conjugates it to isoleucine in *Arabidopsis*. *The Plant Cell* 16: 2117–2127.
- Staswick PE, Su W, Howell SH. 1992. Methyl jasmonate inhibition of root growth and induction of a leaf protein are decreased in an *Arabidopsis thaliana* mutant. *Proceedings of the National Academy of Sciences, USA* 89: 6837–6840.
- Staswick PE, Tiryaki I, Rowe ML. 2002. Jasmonate response locus JAR1 and several related *Arabidopsis* genes encode enzymes of the firefly luciferase superfamily that show activity on jasmonic, salicylic, and indole-3-acetic acids in an assay for adenylation. *The Plant Cell* 14: 1405–1415.
- Stintzi A, Weber H, Reymond P, Browse J, Farmer EE. 2001. Plant defense in the absence of jasmonic acid: the role of cyclopentenones. *Proceedings of the National Academy of Sciences, USA* 98: 12837–12842.
- Taki N, Sasaki-Sekimoto Y, Obayashi T, Kikuta A, Kobayashi K, Aina T, Yagi K, Sakurai N, Suzuki H, Masuda T *et al.* 2005. 12-Oxophytodienoic acid triggers expression of a distinct set of genes and plays a role in wound-induced gene expression in *Arabidopsis*. *Plant Physiology* 139: 1268–1283.
- Tamaoki M, Nakajima N, Kubo A, Aono M, Matsuyama T, Saji H. 2003. Transcriptome analysis of O-3-exposed *Arabidopsis* reveals that multiple signal pathways act mutually antagonistically to induce gene expression. *Plant Molecular Biology* 53: 443–456.
- Thines B, Katsir L, Melotto M, Niu Y, Mandaokar A, Liu G, Nomura K, He SY, Howe GA, Browse J. 2007. JAZ repressor proteins are targets of the SCF<sup>COI1</sup> complex during jasmonate signalling. *Nature* 448: 661–665.
- Thomma BPHJ, Eggermont K, Penninckx IAM, Mauch-Mani B, Vogelsang R, Cammue BPA, Broekaert WF. 1998. Separate jasmonate-dependent and salicylate-dependent defense-response pathways in *Arabidopsis* are essential for resistance to distinct microbial pathogens. *Proceedings of the National Academy of Sciences, USA* 95: 15107–15111.
- Tuominen H, Overmyer K, Keinänen M, Kollist H, Kangasjärvi J. 2004. Mutual antagonism of ethylene and jasmonic acid regulates ozone-induced spreading cell death in *Arabidopsis*. *Plant Journal* 39: 59–69.
- Vahala J, Ruonala R, Keinänen M, Tuominen H, Kangasjärvi J. 2003. Ethylene insensitivity modulates ozone-induced cell death in birch. *Plant Physiology* 132: 185–195.
- Vijayan P, Shockey J, Levesque CA, Cook RJ, Browse J. 1998. A role for jasmonate in pathogen defense of *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* 95: 7209–7214.
- Warner SAJ, Scott O, Draper J. 1992. Characterisation of a wound-induced transcript from the monocotyledon asparagus that shares homology with a new class of pathogenesis-related (PR) proteins. *Plant Molecular Biology* 19: 555–561.
- Wasternack C. 2006. Oxilipins: biosynthesis, signal transduction and action. In: Hedden P, Thomas S. eds. *Plant hormone signaling. Annual plant reviews*. Oxford, UK: Blackwell Publishing Ltd, 185–228.
- Wasternack C. 2007. Jasmonates: an update on biosynthesis, signal transduction and action in plant stress response, growth and development. *Annals of Botany* 100: 681–2007.
- Xie DX, Feys BF, James S, Nieto-Rostro M, Turner JG. 1998. *COI1-1*: an *Arabidopsis* gene required for jasmonate-regulated defense and fertility. *Science* 280: 1091–1094.
- Xu L, Liu F, Lechner E, Genschik P, Crosby WL, Ma H, Peng W, Huang D, Xie D. 2002. The SCF (COI1) ubiquitin-ligase complexes are required for jasmonate response in *Arabidopsis*. *Plant Cell* 14: 1919–1935.
- Xu Y, Chang PF, Liu D, Narasimhan ML, Raghothama KG, Hasegawa PM, Bressan RA. 1994. Plant defense genes are synergically induced by ethylene and methyl jasmonate. *The Plant Cell* 6: 1077–1085.
- Yan JQ, Wang J, Li T, Hwang JR, Patterson C, Zhang H. 2003. AtCHIP, a U-box-containing E3 ubiquitin ligase, plays a critical role in temperature stress tolerance in *Arabidopsis*. *Plant Physiology* 132: 861–869.
- Yan Y, Stolz S, Chetelat A, Reymond P, Pagni M, Dubugnon L, Farmer EE. 2007. A downstream mediator in the growth repression limb of the jasmonate pathway. *The Plant Cell* 19: 2470–2483.
- Yoshioka K, Kachroo P, Tsui F, Sharma SB, Shah J, Klessig DF. 2001. Environmentally sensitive, SA-dependent defense responses in the *cpr22* mutant of *Arabidopsis*. *Plant Journal* 26: 447–459.

Zimmermann P, Hirsch-Hoffmann M, Hennig L, Gruissem W. 2004. GENEVESTIGATOR. *Arabidopsis* microarray database and analysis toolbox. *Plant Physiology* 136: 2621–2632

### Supporting Information

Additional supporting information may be found in the online version of this article.

**Fig. S1** An unaltered acquired thermotolerance in *npr1-1*, *jar1-1* and *ein2-1*.

**Fig. S2** Assessing methyl jasmonate and ethylene effects on heat shock protein gene expression within publicly available transcriptomic databases.

**Fig. S3** Genes exhibiting the highest upregulation in response to methyl jasmonate treatment and to heat shock. Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.